

Accretion in Gravitationally Contracting Clouds

Kohji TOMISAKA*

Faculty of Education, Niigata University, 8050 Ikarashi-2, Niigata 950-21

E-mail: tomisaka@ed.niigata-u.ac.jp.

(Received 1996 June 7; accepted 1996 August 26)

Abstract

Accretion flow in a contracting magnetized interstellar cloud was studied using magnetohydrodynamical simulations and a nested grid technique. First, the interstellar magnetized cloud experiences a “runaway collapse” phase, in which the central density increases drastically within a finite time scale. Finally, it enters an accretion phase, in which inflowing matter accretes onto a central high-density disk or a new-born star. We found that the accretion rate reaches $(4 - 40) \times c_s^3/G$, where c_s and G represent the isothermal sound speed and the gravitational constant, respectively. This is much larger than the standard accretion rate of $0.975c_s^3/G$ for a hydrostatic isothermal spherical cloud (Shu 1977, AAA19.065.044). Due to the effect of an extra infall velocity achieved in the runaway phase ($\sim 2c_s$), the accretion rate is boosted. This rate declines with time in contrast to Shu’s solution, but keeps $\gtrsim 2.5c_s^3/G$. The observed gas infall rate around proto-stars such as L1551 IRS 5 and HL Tau is also discussed.

Key words: Interstellar: clouds — Accretion — Magnetohydrodynamics — Stars: formation — Interstellar: medium

1. Introduction

The physical processes during the course of star formation has attracted much attention in recent years. Many numerical simulations have been performed in order to study the dynamics of the collapse of interstellar clouds

* Visiting fellow of National Astronomical Observatory.

(Larson 1969; Penston 1969; Norman et al. 1980; Narita et al. 1984; for a review, Boss 1987). Even if a cloud is assumed to be initially uniform, the density gradient grows in a dynamical time scale (a sound crossing time). After that, mainly the central part of the cloud contracts. Since the local free-fall time is proportional to $\propto (G\rho)^{-1/2}$, only contraction in the central dense part proceeds. Thus, the central density increases drastically in a finite time (within a couple of sound crossing times), as long as the equation of state remains isothermal. This is called a “runaway collapse”. It is important that the runaway collapse is not stalled by the centrifugal force (Norman et al. 1984), magnetic force (Scott, Black 1980), or even both (Basu, Mouschovias 1994). Thus, it is concluded that if the cloud begins to collapse, the isothermal cloud continues its “runaway collapse” irrespective of the rotation and the magnetic field strength in the cloud.

This is a story of star formation in a cloud far from any hydrostatic equilibria. There is another model, in which the cloud evolves in a quasi-static fashion driven by the plasma drift (ambipolar diffusion; Nakano 1979, 1982, 1983, 1984; Lizano, Shu 1989; Tomisaka et al. 1990; Fiedler, Mouschovias 1993; Ciolek, Mouschovias 1994; Basu, Mouschovias 1994). The neutral molecules flow across the magnetic field lines due to the self-gravitational force, and the ratio of the gas density to the magnetic field strength in a central region increases with time. As a result, the central density increases quasi-statically. However, after the above-mentioned ratio surpasses a critical value, this quasi-static evolution no longer continues, and the cloud begins dynamical contraction.

Shu (1977) studied this dynamical contraction in an isothermal spherical cloud. He assumed that the contraction begins in a cloud having a hydrostatic density distribution with singularity at the center as $\rho(r) = (2\pi Gr^2)^{-1}$. Inflow begins from the center and it propagates outward. Further, the boundary between the inflowing accretion flow and the static, unperturbed part propagates outwardly as a rarefaction wave front with the sound speed. Therefore, this is called “inside-out collapse.” Shu showed that this gives an accretion rate which depends only on the sound speed c_s (and the gravitational constant G) as $\dot{M}_S = 0.975c_s^3/G$, and is constant irrespective of time. However, this depends upon the initial condition that the accretion begins in a *static* cloud. In contrast, the results of dynamical simulations show that there exists a “runaway collapse” even before the accretion flow begins. In this Letter, we consider the complete evolution from the “runaway collapse” to the “inside-out collapse”. We show that more violent accretion occurs, and that the accretion rate is much higher than that of Shu. In the next section, we describe the model. In section 3, numerical results are given and section 4 is devoted to a discussion.

2. Model and Numerical Method

We consider an axisymmetric hydrostatic cloud. To initiate the cloud to collapse, we add a positive density perturbation to the cloud and study its evolution.

The fragmentation of a cylindrical parent cloud with axial magnetic fields leads to a number of clouds threaded with magnetic field lines and connected with each other. We assume that one cloud is formed per length of l_z . If the cylindrical cloud is long, l_z is determined as the wavelength of the gravitational instability with the maximum growth rate λ_{MGR} . The evolution in this case has been studied by Tomisaka (1995: Paper I, 1996: Paper II). These clouds also experience a runaway collapse. Here, we assume $l_z < \lambda_{\text{MGR}}$. When $l_z \ll \lambda_{\text{MGR}}$, there can exist a hydrostatic configuration in such a cloud. As an initial state, we adopt a hydrostatic cloud formed in the case of $l_z < \lambda_{\text{MGR}}$.

The magnetohydrostatic equilibrium is obtained by solving the coupled equations (Mouschovias 1976; Tomisaka et al. 1988) of the Poisson equation for the gravitational potential and the Grad-Shafranov equation for the magnetic vector potential. We search for solutions by an iterative method, the so-called self-consistent field method. Since this solution is in hydrostatic equilibrium, the structure remains unchanged after time integration. Thus, a positive density perturbation with a relative amplitude of 10% is added to the central several grids.

After choosing the units as ρ_s (the density on the cloud surface) = $c_s = 4\pi G = 1$, the model contains 4 parameters: the plasma beta of the virtual parent cylindrical cloud, β , the center-to-surface density ratio of this parent cloud, F , the length of the cloud, l_z , and the mass of the cloud, M_{cl} . The assumed model parameters are given in table 1.

As boundary conditions, we assume that far from the cloud the magnetic field approaches a uniform one as $(B_z, B_r) = ((8\pi p_{\text{ext}}/\beta)^{1/2}, 0)$, where the external pressure p_{ext} is taken to be equal to the surface pressure $\rho_s c_s^2$. On the outer boundary of a numerical box, the gravitational potential ψ reduces to that made by a series of point masses of M_{cl} , which are separated from each other by l_z on the z -axis.

The basic equations to be solved are the magnetohydrodynamics and Poisson equations in cylindrical symmetry. As a numerical scheme, we employed van Leer's (1977) monotonic interpolation and the constrained transport (Evans, Hawley 1988) for the magnetic induction equation. As a Poisson solver, we adopted the preconditioned conjugate gradient squared method. To resolve the fine structure near the center of the cloud, the nested grid technique was employed (for the detail of this numerical scheme, see Paper II). This nested grid method using 14 levels of grids has an effective resolution corresponding to a simple finite-difference scheme using $10^6 \times 10^6$ grid points.

3. Results

The initial structure of the static cloud is shown in figure 1a for L2 (Level 2 of the nested grid system; L_n has twice finer resolution than L_{n-1}). First, the perturbation grows in proportion to $\delta\rho \propto \exp(\omega t)$, which is characteristic to linear growth. Since a flow crossing the magnetic field lines is blocked by the field, a disk which runs perpendicular to the magnetic fields is formed. After $t \sim 0.18$, the central density increases rapidly with time, and at $t = 0.2115$ the density reaches $10^{10}\rho_s$ in the runaway collapse phase (the evolution in this phase was described previously in Papers I and II). Even at this time, the global structure is unchanged from that shown in figure 1a. The structure of the central region (L7) at this stage is shown in figure 1b. Gas is contracting onto the disk flowing along the magnetic field lines, which run perpendicular to the disk, with $|v_z| \gg |v_r|$. Outward-facing shock waves, whose fronts are parallel to the disk, are formed. The magnetic field lines are pulled radially inwardly in the disk where the thermal pressure dominates the magnetic one. The cloud now enters the accretion phase.

To continue the simulation, we adopted so-called “sink cell method,” in which the matter in the grids near the center is assumed to accrete onto the central high-density body, and to be removed from the cells (Boss, Black 1982). We assume that the sink cells distribute inside a distance of 16-times the grid size from the center in L7 and only in the sink cells in which $\rho > \rho_{\text{sink}}$, the excess density ($\rho - \rho_{\text{sink}}$) and the corresponding momentum are both removed (here we take $\rho_{\text{sink}} = 10^6\rho_s$). The removed mass in each time steps accounts for the increase in the mass of the central body. As a result of the second assumption, sink actually occurs in a small number of central grids. Compared with a reference calculation with a different size of the sink region, it is confirmed that the size of the sink region is not important to the result.

The structure at $\tau = 0.0569$ after the beginning of accretion is shown in figure 1c. (In this Letter, t and τ represent the time from the perturbation was given and that from the accretion began.) It clearly shows that the magnetic fields are squeezed, and run almost radially near the disk. The shape of the magnetic field lines becomes like an hourglass. Comparing figures 1b and 1c, it is shown that the disk continues to contract, and becomes thinner during the accretion phase. From figure 1d, it is shown that the contracting speed exceeds the sound speed and reaches $\simeq 5c_s$ (near $r \simeq 0.02$), while in the runaway collapse phase it is not more than $(1.5\text{-}2)c_s$.

The accreted mass is calculated as the mass removed from the sink cells. In figure 2, we illustrate the time evolution of the accretion rate $\dot{M}$ normalized to c_s^3/G by the thick lines. This clearly shows that the rate is *not*

constant with time. Since the value for Shu's solution is equal to 0.975, the difference attains $\gtrsim 40$ times just after accretion begins. Even in the late phase, the accretion rate is $\gtrsim 2.5$ -times larger than that expected in a static cloud (0.975). The time-averaged accretion rate ($\int_0^\tau \dot{M} dt/\tau$) is equal to $\simeq 4c_s^3/G$. This is explained by the extra inflow that the cloud achieved in the runaway phase. Gas within $r \lesssim 0.02$ is falling at a speed of $|v_r| \gtrsim 1.5c_s$ (figure 1b) when accretion begins. This enhances the accretion rate by an amount $\sim \rho|v_r|4\pi R^2$ if spherical symmetry is assumed.

To see the effect of the initial condition on the accretion rate, we studied model B, which has a lower density contrast (ρ_c/ρ_s) and a larger mass (M_{cl}) than those of model A. In this case, a density perturbation with a relative amplitude of 20% is necessary for the cloud to begin contraction, while this is 10% for model A. This means that the initial condition of model B is much more stable than that of model A for gravitational contraction. However, model B indicates a larger accretion rate by contraries (thin lines of figure 2). It seems to result from the fact that the runaway collapse phase lasts for 0.96 in contrast to 0.21 (model A). This indicates that the extra inflow is considerably accelerated in the runaway phase, and that the inflow achieved in the runaway phase plays an important role concerning the later accretion rate.

4. Discussion

A high accretion rate has been argued by Whitworth and Summers (1985) as well as Foster and Chevalier (1993) for spherical symmetric clouds. The former studied various *self-similar* spherically symmetric solutions, and found that the accretion rate reaches $46.84c_s^3/G$ in the case that the accretion begins in the Larson(1969) and Penston (1969) self-similar solution at $\tau = 0$ (The Larson and Penston self-similar solution itself has a constant mass inflow rate as $29c_s^3/G$). The Larson and Penston self-similar solution is believed to be attained near the center of the cloud in a runaway (dynamical) contraction which occurs in an initially diffuse cloud. Thus, it seems natural that in the beginning of the accretion phase the accretion is very active as $\dot{M} \simeq (40-50) c_s^3/G$, as shown in our MHD simulations. Foster and Chevalier (1993) confirmed this by their one-dimensional hydro-code, and studied cloud contraction from $\tau < 0$ to $\tau > 0$ throughout. They found that the accretion rate attains $\sim 10c_s^3/G \gg \dot{M}_S$, and then gradually declines.

The accretion phase of a magnetized cloud has been studied by Galli and Shu (1993a, b). They showed that the accretion rate is not affected very much by the effect of magnetic fields, e.g., $\dot{M} \simeq 0.975c_s^3/G$; this comes from the

subtle role of magnetic fields in the dynamics. The latter fact is also confirmed in this Letter, since the accretion rate is not largely different from that calculated for non-magnetic clouds (Whitworth, Summers 1985). However, the accretion rate of Galli and Shu is much smaller than ours. This is explained as follows. They considered a cloud such as a hydrostatic singular isothermal sphere threaded by uniform magnetic fields, i.e., accretion is assumed to begin in this static cloud. In contrast, since an extra infall velocity is achieved during the dynamical runaway collapse, our model indicates a larger accretion rate than does theirs.

The normalization adopted here gives the actual physical scales: for time $t = (4\pi G\rho_s)^{-1/2} = 1.75(n_s/100\text{ cm}^{-3})^{-1/2}\text{ Myr}$, where the n_s represents the particle density at the surface of the cloud, for an accretion rate $\dot{M} = c_s^3/G = 1.62 \times 10^{-6}(c_s/190\text{ m s}^{-1})^3 M_\odot\text{ yr}^{-1}$, and for mass $M = \dot{M}t = c_s^3/G/(4\pi G\rho_s)^{1/2} = 2.84(c_s/190\text{ m s}^{-1})^3(n_s/100\text{ cm}^{-3})^{-1/2}M_\odot$. Here, we assumed that the temperature is equal to 10 K. Thus, if we choose the above normalization, it is shown that a mass of $M = 1.78 M_\odot$ has accreted in $\tau = 0.21\text{ Myr}$ (final state of figure 2). The average accretion rate becomes $8.7 \times 10^{-6}M_\odot\text{ yr}^{-1}$. In contrast, the model by Shu (1977) predicts $1.58 \times 10^{-6}M_\odot\text{ yr}^{-1}$ for the standard $A = 2$ model and $9.04 \times 10^{-6}M_\odot\text{ yr}^{-1}$ for the $A = 4$ model. [This A is a density parameter to specify the initial density distribution as $\rho(r) = Ac_s^2/(4\pi G)r^{-2}$. The distribution coincides with that of the singular hydrostatic solution (Chandrasekhar 1957) only when $A = 2$. When $A > 2$, the solution is out of hydrostatic balance and the cloud is contracting. Since accretion occurs in a dynamically-contracting cloud core in our model, it is reasonable that the accretion rate is similar to that expected by the Shu's model with $A = 4$.]

It is not easy to determine the accretion rate from the direct detection of infalling gas. However, there exist a number of observations. From a combination of the infall velocity (0.55 km s^{-1}) and the number density ($2 \times 10^5\text{ cm}^{-3}$) at a radius from the center $20''$ of L1551 IRS 5, Fuller et al. (1995) obtained a mass accretion rate to this object as $\dot{M} \lesssim 1.4 \times 10^{-5}M_\odot\text{ yr}^{-1}$. This is rather larger than Shu's accretion rate mentioned above. However, this can be naturally explained by our model. Applying our models to fit this accretion rate, the age should be $3 \times 10^4\text{ yr}$ (model A) or $8 \times 10^4\text{ yr}$ (model B) for this object. This is consistent with the age determined from the dynamical age of the outflow, $3 \times 10^4\text{ yr} - 10^5\text{ yr}$ (Moriarty-Schieven, Snell 1988). Another example is HL Tau. From the Nobeyama Millimeter Array map, Hayashi et al. (1993) estimated the mass infall rate at 1400 AU from the center as $5 \times 10^{-6}M_\odot\text{ yr}^{-1} - 2 \times 10^{-5}M_\odot\text{ yr}^{-1}$. Since the accretion rate is proportional to c_s^3 , a large accretion rate may be explained by a slightly higher temperature, such as 20 K. However, these accretion rates are naturally explained by our model without assuming a higher temperature. New observations with a finer spatial resolution are desired in

order to resolve the inflow velocity profile and to determine the dynamics in the contracting cloud further.

The work was supported in part by Grant-in-Aids for Scientific Research from Ministry of Education, Science, Sports and Culture (07640351, 07304025).

References

- Basu S., Mouschovias T. Ch. 1994, ApJ 432, 720
- Boss A. P. 1987, in *Interstellar Processes* (Reidel: Dordrecht), ed D. J. Hollenbach, H. A. Thronson, Jr. p.321
- Boss A. P., Black D. C. 1982, ApJ 258, 270
- Chandrasekhar S. 1939, *An Introduction to the Study of Stellar Structure* (University Chicago Press: Chicago), section 22
- Ciolek G. E., Mouschovias T. Ch. 1994, ApJ 425, 142
- Evans C. R., Hawley J. F. 1988, ApJ 332, 659
- Fiedler R. A., Mouschovias T. Ch. 1993, ApJ 415, 680
- Foster P. N., Chevalier R. A. 1993, ApJ 416, 303
- Fuller G. A., Ladd E. F., Padman R., Myers P. C., Adams F. C. 1995, ApJ 454, 862
- Galli D., Shu F. H. 1993a, ApJ 417, 220
- Galli D., Shu F. H. 1993b, ApJ 417, 243
- Hayashi M., Ohashi N., Miyama S. M. 1993, ApJ 418, L71
- Larson R. B. 1969, MNRAS 145, 271
- Lizano S., Shu F. H. 1989, ApJ 342, 834
- Moriarty-Schieven G. H., Snell R. L. 1988, ApJ 332, 364
- Mouschovias T. Ch. 1976, ApJ 206, 753
- Nakano T. 1979, PASJ 31, 697
- Nakano T. 1982, PASJ 34, 337
- Nakano T. 1983, PASJ 35, 87
- Nakano T. 1984, *Fund. Cosmic Phys.* 9, 139
- Narita S., Hayashi C., Miyama S. M. 1984, *Prog. Theor. Phys.* 72, 1118
- Norman M. L., Wilson J. R., Barton R. T. 1980, ApJ 239, 968
- Penston M. V. 1969, MNRAS 144, 425
- Scott E. H., Black D. C. 1980, ApJ 239, 166

Shu F. H. 1977, ApJ 214, 488

Tomisaka K. 1995, ApJ, 438, 226 (Paper I)

Tomisaka K. 1996, PASJ, 48, No.5 (Paper II)

Tomisaka K., Ikeuchi S., Nakamura T. 1988, ApJ 326, 208

Tomisaka K., Ikeuchi S., Nakamura T. 1990, ApJ 362, 202

van Leer B. 1977, J. Comp. Phys. 23, 276

Whitworth A., Summers D. 1985, MNRAS 214, 1

Table 1. Model parameters.

Model	β^*	$F^\dagger$	$\rho_c/\rho_s^{\ddagger}$	$M_{\text{cl}}^\S$	$l_z^\P$
A.....	1	100	1000	37.96	12.73
B.....	1	100	100	46.4	12.73

*The plasma β in the parent cylindrical cloud.

† The center-to-surface density ratio in the parent cloud.

‡ The center-to-surface density ratio of the hydrostatic cloud.

§ The mass of the cloud.

¶ The length of the parent cloud.

Figure Captions

Fig. 1. (a) Initial state ($t = 0$). Density contour lines (closed lines) and magnetic field lines (running horizontally) are plotted for the L2 grid. The steps of the density contours are taken as $\rho = 10^{3n/20}$ ($n = 0, 1, \dots, 20$). The z -axis, the symmetry axis, is taken horizontally. (b) The structure at the end of the runaway collapse phase ($t = 0.2115$) in the L7 grid. A disk is formed running perpendicular to the global magnetic field lines. The size of the grid is $2^5 = 32$ times finer than that of (a). (c) The structure in the accretion phase at $t = 0.2684$ ($\tau = 0.0569$). The size of the grid is the same as (b). (d) Radial distributions of the density (ρ : solid line), the radial velocity (v_r : dotted line) and the magnetic field strength (B_z : dashed line) on the equatorial plane ($z = 0$).

Fig. 2. Mass accretion rate against the age after the core formation τ . The rate is normalized to c_s^3/G . The solid and dotted lines represent the instantaneous accretion rate as $\dot{M}(\tau)$ and the time averaged one as $\int_0^\tau \dot{M}(t)dt/\tau$, respectively. The thick and thin lines correspond to models A and B, respectively. The value obtained by Shu (1977) is 0.975. On the upper and right axes, typical dimensional values are plotted with assuming the conversion factors in section 4 as $t = (4\pi G \rho_s)^{-1/2} = 1.75(n_s/100 \text{ cm}^{-3})^{-1/2} \text{ Myr}$ and $\dot{M} = c_s^3/G = 1.62 \times 10^{-6} (c_s/190 \text{ m s}^{-1})^3 M_\odot \text{ yr}^{-1}$.